



WWJMRD2023; 9(07):108-110  
www.wwjmr.com  
International Journal  
Peer Reviewed Journal  
Refereed Journal  
Indexed Journal  
Impact Factor SJIF 2017:  
5.182 2018: 5.51, (ISI) 2020-  
2021: 1.361  
E-ISSN: 2454-6615

**Noviani Prasetyaningsih**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Anggraeni Adiwardhani**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Riani Witjaksana**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Isa Bella**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Jihan Samira**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Ida Effendi**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

**Monica Dwi Hartanti**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia  
Center for Biomedical  
Research, National Research  
and Innovation Agency, West  
Java, Indonesia.

**Freily Akay**  
Universitas Sam Ratulangi,  
Manado, Indonesia.

**Correspondence:**  
**Monica Dwi Hartanti**  
Faculty of Medicine,  
Universitas Trisakti, Jakarta,  
Indonesia.

## Dry eye disease and its risk factors in rural and urban areas in Indonesia

**Noviani Prasetyaningsih, Anggraeni Adiwardhani, Riani Witjaksana, Isa Bella, Jihan Samira, Ida Effendi, Monica Dwi Hartanti, Freily Akay**

### Abstract

This cross-sectional study was conducted in two regions in Indonesia, Karawaci Banten and Depok City to determine the risk factors of Dry Eye Disease (DED) in rural and urban areas. To assess DED, the Schirmer test was performed on one eye (right eye). The risk factors being studied were age, gender, smoking habits, and the use of electronic devices. A total of 145 respondents was recruited, with 60 respondents (41.4%) from rural areas and 85 respondents (58.6%) from urban areas. The prevalence of DED was higher in urban areas, with 54.1% (46/85) compared to 48.3% (29/60) in rural areas. There was no significant association between age and gender with the incidence of DED in both rural and urban areas ( $p > 0.05$ ). Smoking was significantly associated with DED incidence in rural areas ( $p = 0.0004$ ), but not in urban areas. The use of electronic devices was significantly associated with DED incidence only in urban areas ( $p = 0.043$ ). Smoking is a risk factor for DED in rural areas, while the use of electronic devices is the main risk factor in urban areas.

**Keywords:** Dry Eye Disease, Rural, Urban, Schirmer.

### 1. Introduction

Dry eye disease is defined by the Report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop as a multifactorial disease of the tears and ocular surface, which results in symptoms of discomfort, visual disturbance, and tear film instability, with potential damage to the ocular surface.<sup>[1]</sup> The prevalence of dry eye disease ranges from 5-35% of the population worldwide, with Nigeria having a prevalence of 19.2%,<sup>[2]</sup> China 17%,<sup>[3]</sup> and Dubai 62.6%.<sup>[4]</sup> In Indonesia, the prevalence of dry eye disease is 27.5% in a study conducted in the Riau Islands.<sup>[5]</sup> The purpose of this study is to compare the incidence of dry eye disease in rural and urban areas based on risk factors such as age, gender, smoking habits and the use of electronic devices.

### 2. Methods

This cross-sectional study was conducted in two regions in Indonesia, Karawaci Banten as a rural area and Depok City as an urban area. The exclusion criteria were the presence of signs of eye infection, the use of contact lenses, and the use of routine eye drops for the last 6 months. After signing the informed consent and filling out the identity sheet, there were several additional questions regarding smoking habits, and the use of electronic devices. The eye examination performed was a Schirmer's test just to the right eye to assess the quantity of tears. The Schirmer test was performed by an ophthalmologist without anesthetic using Whatman No. 41 filter paper, which is 5 mm wide and 30 mm long. The tip of the Schirmer paper was folded about 5 mm and inserted into the inferior fornix conjunctiva. The eye was closed and left for 5 minutes. After 5 minutes, the length of the wetted Schirmer paper was measured. DED was indicated if the wetted Schirmer paper was less than 10 mm.<sup>[6]</sup>

All data were analyzed using the GraphPad version 9 program. The data collected was presented in the form of frequency and percentage distributions. Bivariate analysis of two groups was performed using the Chi-square test, if certain conditions were met with a significance level of  $p < 0.05$ . If not, the Fisher's exact test was used.

**2. Result**

A total of 145 respondents participated in the research, with 60 respondents (41.4%) from rural areas and 85 respondents (58.6%) from urban areas. The respondents were divided into two age groups, the age group of 30-50 years old consisted of 79 respondents (54,5%), while the age group of over 50 years old consisted of 66 respondents (45,5%). The prevalence of DED was 48.3% in the rural group (n=60) and 54.1% in the urban group (n=85). The demographic profile of the study subjects in rural and urban areas is presented in Table 1.

Table 2 shows various risk factors for Dry Eye Disease in both rural and urban areas. According to our research, age and gender did not have a significant relationship with the incidence of Dry Eye Disease ( $p > 0.05$ ). However, smoking habits were significantly associated with the incidence of Dry Eye Disease in rural areas ( $p = 0.0004$ ), but not in urban areas. On the other hand, excessive use of electronic devices was the main risk factor for Dry Eye Disease in urban areas ( $p = 0.043$ ).

**Table 1:** Characteristics of respondents in Rural and Urban areas.

Characteristics	Rural		Urban	
	n = 60	%	n = 85	%
Gender				
Male	26	43,3	33	38,8
Female	34	56,7	52	61,2
Ages				
30 - 50	44	73,3	35	41,2
More than 50	16	26,7	50	58,8
Smoking habits				
Yes	13	21,7	23	27,1
No	47	78,3	62	72,9
Use of electronic devices				
Yes	43	71,7	33	38,8
No	17	28,3	52	61,2
Dry Eye (Schirmer)				
Yes	29	48,3	46	54,1
No	31	51,7	39	45,9

**Table 2:** Distribution of Dry Eye Disease based on Risk Factors in Rural and Urban Areas.

Risk Factors	Rural (n = 60)				p	Urban (n = 85)				p
	Dry Eye (Schirmer)					Dry Eye (Schirmer)				
	Yes		No			Yes		No		
	n	%	n	%		n	%	n	%	
Gender										
Male	15	25,0	11	18,3	0,205	14	16,5	19	22,4	0,085
Female	14	23,3	20	33,3		32	37,6	20	23,5	
Ages										
30 - 50	19	31,7	25	41,7	0,185	19	22,4	16	18,8	0,979
More than 50	10	16,7	6	10,0		27	31,8	23	27,0	
Smoking habits										
Yes	12	20,0	1	1,7	0,0004*	9	10,6	15	17,6	0,054
No	17	28,3	30	50,0		37	43,5	24	28,2	
Use of electronic devices										
Yes	21	35,0	22	36,7	0,901	17	20,0	23	27,1	0,043
No	8	13,3	9	15,0		29	34,1	16	18,8	

\* Fisher test

**4. Discussion**

The prevalence of DED based on Schirmer examination in this study was 48.3% in rural areas and 54.1% in urban areas. This number has increased compared to the results of a study conducted in Indonesia in 2002, which had a prevalence of 27.5%.<sup>[5]</sup> A relatively high prevalence was also found in Hotan, China, where the prevalence reached 40.6%,<sup>[7]</sup> while the prevalence in a study in rural areas of India was 45.39%.<sup>[8]</sup> A study in Ghana, Africa compared the prevalence of DED in rural and urban areas, and the results showed that respondents in rural areas suffered more from DED and had more severe symptoms than respondents in urban areas.<sup>[9]</sup> DED usually affects older individuals, with its prevalence increasing as age increases. Women are at a higher risk of developing DED compared

to men.<sup>[10-15]</sup> This is supported by clinical examinations that have found a decrease in tear production in individuals in their sixth decade of life.<sup>[10]</sup>

In his research on the effects of smoking on tear proteins, Grus found that there were changes in the tear film of a smoker, where there was an increase in protein spots due to toxins from cigarettes that caused oxidative damage to the proteins.<sup>[16]</sup> This explains why smoking is a risk factor for Dry Eye Disease. The research conducted in Saudi Arabia regarding electronic cigarettes, which are claimed to be an alternative to conventional cigarettes that are safe and non-addictive, has found that they are actually unsafe and can cause health issues such as nausea, vomiting, headaches, dryness around the mouth, eyes, and mucous membranes.<sup>[17]</sup> In our research, there is a positive

correlation between smoking habits and the incidence of Dry Eye Disease in rural areas ( $p=0,0004$ ).

Several literature states that excessive use of electronic devices can cause dry eyes. A study in Serbia with a population of 18–34-year-old students found a prevalence of Dry Eye Disease (DED) of 60.5%. Electronic devices have a significant relationship with the occurrence of DED, and the longer the duration of electronic device use, the higher the prevalence of DED. [18] To assess the effect of smartphones on the onset of DED symptoms, tear film and oxidative stress in the eyes, Choi conducted a case-control study, and the results showed that using electronic devices for more than 4 hours would worsen DED symptoms and lead to oxidative stress in the eyes.

## 5. Conclusion

The habit of smoking and excessive use of electronic devices can cause damage to the tear film layer, resulting in DED symptoms. It is necessary to explain this to the public, both in rural and urban areas.

## References

1. International Dry Eye WorkShop. The definition and classification of dry eye disease: report of the Definition and Classification Subcommittee of the International Dry Eye WorkShop. *Ocul Surf.* 2007;5:75-92.
2. Echieh CI, Etim BA, Echieh CP, Oyeniyi T, Ajewole J. A comparative assessment of Dry Eye Disease Among Outdoor Street Sweepers and Indoor Office Cleaners. *BMC Ophthalmology* 2021;21:265. doi.org/10.1186/s12886-021-02025-y
3. Liu N, Liu L, Li J, Sun Y. Prevalence of and Risk Factors for Dry Eye Symptom in Mainland China: A Systematic Review and Meta-Analysis. *J. Ophthalmol* 2014;1-8. http://dx.doi.org/10.1155/2014/748654
4. Alkabbani S, Jeyaseelan L, Rao AP, Thakur SP, Warhekar PT. The Prevalence, Secerity, and Risk Factors for Dry Eye Disease in Dubai – a cross sectional study. *BMC Ophthalmology* 2021;21:219. https://doi.org/10.1186/s12886-021-01978-4
5. Lee AJ, Lee J, Saw S-M, Gazzard G, Koh D, Widjaja D, Tan DTH. Prevalence and Risk Factors Associated with Dry Eye Symptoms: a population-based study in Indonesia. *Br J Ophthalmol* 2002;86:1347–1351
6. Stevens S. Schirmer's test. *Community Eye Health.* 2011 Dec;24(76):45
7. Li X, Wang Z, Mu J, Puerkaiti H, Nulahou A et al. Prevalence and Associated Risk Factors of Dry Eye Disease in Hotan, Xinjiang: A Cross Sectional Study. *BMC Ophthalmology* (2023); 23:214. https://doi.org/10.1186/s12886-023-02955-9
8. Ranjan R, Shukla SK, Singh CV, Mishra BN, Sinha S et al. Prevalence of Dry Eye and Its Association with Various Risk Factors in Rural Setup of Western Uttar Pradesh in a Tertiary Care Hospital. *Journal of Preventive Medicine,* 6, 57-63. http://dx.doi.org/10.4236/ojpm.2016.61005\_
9. Osaie EA, Ablordeppey RK, Horstmann J, Kumah DB, Steven P. Clinical Dry Eye and Meibomian Gland Features Among Dry Eye Patients in Rural and Urban Ghana. *Clinical Ophthalmology* 2020;14:4055-63
10. de Paiva CS. Effects of Aging in Dry Eye. *Int Ophthalmol Clin.* 2017; 57(2): 47–64. doi:10.1097/HIO.000000000000170
11. Ozdemir M, Temizdemir H. Age and gender related tear function changes in normal population. 2010. *Eye* 24; 79-83
12. Sharma A, Hindman HB. Aging: A Predisposition to Dry Eyes. 2014. *Journal of Ophthalmology.* http://dx.doi.org/10.1155/2014/781683
13. Bikbov MM, Kazakbaeva GM, Rakhimova EM, Rusakova IA, Fakhretdinova AA et al. The prevalence of dry eye in a very old population. *Acta Ophthalmologica.* 2022; 100: 262-8. doi: 10.1111/aos.14937
14. AlMarshedi MM, Alshammari SA. The Prevalence of Dry Eye Disease and Related Factors Among Adult Patients Attending Primary Healthcare Centers in Riyadh, Saudi Arabia. *Cureus* 14(11): e31400. DOI 10.7759/cureus.31400
15. Panggat KMS, Covar RV SL. Prevalence of Dry Eye Disease in an Urban Community. *Philipp J Ophthalmol.* 2015;40:29-35
16. Grus FH, Sabuncuo P, Augustin A, Pfeiffer N. Effect of Smoking on Tear Protein. *Graefe's Arch Clin Exp Ophthalmol.* (2002) 240:889–892. DOI 10.1007/s00417-002-0539-y
17. Meo SA, Al Asiri SA. Effects of Electronic Cigarette Smoking on Human Health. *European Review for Medical and Pharmacological Sciences.* 2014; 18: 3315-3319
18. Aćimović L, Stanojlović S, Kalezić T, Krnjaja BD. Evaluation of Dry Eye Symptoms and Risk Factors among Medical Students in Serbia. *Plos One* 2022;17(10): e0275624. https://doi.org/10.1371/journal.pone.0275624
19. Wu H, Wang Y, Dong N, Yang F, Lin Z et al. Meibomian Gland Dysfunction Determines the Severity of the Dry Eye Conditions in Visual Display Terminal Workers. *Plos One* 2014;9(8): e105575. doi:10.1371/journal.pone.0105575
20. Choi JH, Li Y, Kim SH, Jin R, Kim YH et al. The influences of smartphone use on the status of the tear film and ocular surface. *Plos One.* 2018;13(10): e0206541. https://doi.org/10.1371/journal.pone.0206541